

# AUROQUIL COLD AND FLU NIGHTTIME RELIEF - acetaminophen, doxylamine succinate, and dextromethorphan hbr solution

Aurohealth LLC

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## Drug Facts

Active ingredients (in each 30 mL dose cup)	Purpose
Acetaminophen, USP 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr, USP 30 mg	Cough suppressant
Doxylamine succinate, USP 12.5 mg	Antihistamine

## Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

## Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- use dose cup or tablespoon (TBSP)
- do not exceed 4 doses (120 mL) (8 TBSP) per 24 hours
- mL = milliliter; TBSP = tablespoon

adults & children 12 years & over	30 mL (2 TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

- **When using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

#### **Other information**

- **each 30 mL dose cup contains:** potassium 5 mg, sodium 37 mg
- store at 20° to 25°C (68° to 77°F)

#### **Inactive ingredients**

acesulfame potassium, alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

#### **Questions or Comments?**

**1-855-274-4122**

Distributed by:

**Aurohealth LLC.**

2572 Brunswick Pike,

Lawrenceville, NJ 08648

#### **PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 6 FL OZ (177 mL Bottle)**

#### **AUROHEALTH**

Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*

**NDC 58602-137-18**

**AuroQuil**

**COLD & FLU**

**Nighttime Relief**

Each dose (per 30 mL) (2 TBSP) of oral solution contains:

650 mg - **Acetaminophen** USP (Pain reliever/fever reducer)

12.5 mg - Doxylamine succinate USP (Antihistamine)

30 mg - Dextromethorphan HBr USP (Cough suppressant)

- **Aches, Fever & Sore Throat**
- **Sneezing, Runny Nose**
- **Cough**

**Alcohol 10%**

**6 FL OZ (177 mL)**

# Front

**Tamper Evident:** Do not use if printed shrinkband is missing or broken.  
Failure to follow these warnings could result in serious consequences

3001000854



\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu

Distributed by:  
**Aurohealth LLC.**  
2572 Brunswick Pike,  
Lawrenceville, NJ 08648

## PARENTS:

Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

LOT:

EXP:



Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*  
NDC 58602-137-18

## AuroQuil

**COLD & FLU**

### Nighttime Relief

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - Acetaminophen USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough suppressant)

- ◆ Aches, Fever & Sore Throat
- ◆ Sneezing, Runny Nose
- ◆ Cough

Alcohol 10%

6 FL OZ (177 mL)

Lift Here for continued **Drug Facts**

## Drug Facts

### Active ingredients (in each 30 mL dose cup)

Acetaminophen, USP 650 mg...Pain reliever/fever reducer  
Dextromethorphan HBr, USP 30 mg...Cough suppressant  
Doxylamine succinate, USP 12.5 mg...Antihistamine

### Purpose

**Uses** temporarily relieves common cold/flu symptoms:  
■ cough due to minor throat and bronchial irritation  
■ sore throat ■ headache ■ minor aches and pains  
■ fever ■ runny nose and sneezing

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening

# Back

## Drug Facts (continued)

■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleep

## Drug Facts (continued)

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland ■ a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

**When using this product** ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

# Bottom Ply

## Drug Facts (continued)

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed - see Overdose warning
- use dose cup or tablespoon (TBSP)

## Drug Facts (continued)

■ do not exceed 4 doses (120 mL) (8 TBSP) per 24 hours ■ mL = milliliter; TBSP = tablespoon  
adults & children 12 years & over 30 mL (2 TBSP) every 6 hours  
children 4 to under 12 years ask a doctor  
children under 4 years do not use

■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

**Other information** ■ each 30 mL dose cup contains: potassium 5 mg, sodium 37 mg ■ store at 20° to 25°C (68° to 77°F)

### Inactive ingredients

acesulfame potassium, alcohol, anhydrous citric acid, D&C yellow no.10, FD&C green no.3, FD&C yellow no.6, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or Comments?** © 1-855-274-4122

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 8 FL OZ (237 mL Bottle)**

**AUROHEALTH**

Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*



NDC 58602-137-20

**AuroQuil**  
**COLD & FLU**  
**Nighttime Relief**

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - **Acetaminophen** USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough suppressant)

- **Aches, Fever & Sore Throat**
- **Sneezing, Runny Nose**
- **Cough**

**Alcohol 10%**

**8 FL OZ (237 mL)**

## Front

3001000355

**Tamper Evident: Do not use if printed shrinkband is missing or broken. Failure to follow these warnings could result in serious consequences**



\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu

Distributed by: **Aurohealth LLC.**  
2572 Brunswick Pike,  
Lawrenceville, NJ 08648

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

LOT:  
EXP:

**AUROHEALTH**

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu\*  
NDC 58602-137-20

# AuroQuil

## COLD & FLU

### Nighttime Relief

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - Acetaminophen USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough suppressant)

- ◆ Aches, Fever & Sore Throat
- ◆ Sneezing, Runny Nose
- ◆ Cough

**Alcohol 10%** **8 FL OZ (237 mL)**

Lift Here for continued **Drug Facts**

Active ingredients (in each 30 mL dose cup)	Purpose
Acetaminophen, USP 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr, USP 30 mg	Cough suppressant
Doxylamine succinate, USP 12.5 mg	Antihistamine

**Uses**  
temporarily relieves common cold/flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while

## Back

**Drug Facts** (continued)

using this product.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

**Drug Facts** (continued)

contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleep

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma  
■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema  
■ trouble urinating due to enlarged prostate gland ■ a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers  
■ taking the blood thinning drug warfarin

**When using this product** ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug

may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

## Bottom Ply

### Drug Facts (continued)

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

■ take only as directed - see Overdose warning  
■ use dose cup or tablespoon (TBSP) ■ do not exceed 4 doses (120 mL) (8 TBSP) per 24 hours  
■ mL = milliliter; TBSP = tablespoon

### Drug Facts (continued)

adults & children 12 years & over	30 mL (2 TBSP) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

### Other information

■ each 30 mL dose cup contains: potassium 5 mg, sodium 37 mg  
■ store at 20° to 25°C (68° to 77°F)

### Inactive ingredients

acesulfame potassium, alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or Comments?** © 1-855-274-4122

## PACKAGE LABEL-PRINCIPAL DISPLAY PANEL 12 FL OZ (355 mL Bottle)

### AUROHEALTH

Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*

NDC 58602-137-16

AuroQuil

COLD & FLU

Nighttime Relief

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - **Acetaminophen** USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough suppressant)

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose
- Cough

Alcohol 10%

12 FL OZ (355 mL)



Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*

NDC 58602-137-16

# AuroQuil

## COLD & FLU

### Nighttime Relief

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - Acetaminophen USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough suppressant)

- ◆ Aches, Fever & Sore Throat
- ◆ Sneezing, Runny Nose
- ◆ Cough

Alcohol 10% 12 FL OZ (355 mL)

**Purpose**

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine

**Drug Facts**

**Active Ingredients (in each 30 mL dose cup)**

Acetaminophen, USP 650 mg  
Dextromethorphan HBr, USP 30 mg  
Doxylamine succinate, USP 12.5 mg

**Uses** temporarily relieves common cold/flu symptoms: cough due to minor throat and bronchial irritation sore throat minor aches and pains fever runny nose and sneezing

**Warnings** Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash if a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Do not make a child sleep. Ask a doctor before use if you have liver disease glaucoma cough that occurs with too much phlegm (mucus) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema trouble urinating due to enlarged prostate gland a sodium-restricted diet Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers taking the blood thinning drug warfarin When using this product do not use more than directed excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** Take only as directed - see Overdose warning use dose cup or tablespoon (TBSP) do not exceed 4 doses (120 mL) (8 TBSP) per 24 hours  
ml = milliliter; TBSP = tablespoon  
when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

**Other information** each 30 mL dose cup contains: potassium 5 mg, sodium 37 mg store at 20° to 25°C (68° to 77°F)

**Inactive ingredients** acetic acid, potassium, alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C yellow no. 6, flavors, high fructose corn syrup, polyethylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or Comments?** 1-855-274-4122

**Tamper Evident:** Do not use if printed shrinkband is missing or broken. Failure to follow these warnings could result in serious consequences

**PARENTS:** 3001000356

Distributed by:  
AuroHealth LLC  
2572 Brunsview Pike  
Lawrenceville, GA 30046

3001000356

58602173716

## PACKAGE LABEL-PRINCIPAL DISPLAY PANEL TWO 12 FL OZ (355 mL Bottles)

VALUE  
Twin Pack

### AUROHEALTH

Compare to the active ingredients  
in Vicks® NyQuil® Cold & Flu\*

NDC 58602-137-35

### AuroQuil COLD & FLU Nighttime Relief

Each dose (per 30 mL) (2 TBSP) of oral solution contains:  
650 mg - **Acetaminophen** USP (Pain reliever/fever reducer)  
12.5 mg - Doxylamine succinate USP (Antihistamine)  
30 mg - Dextromethorphan HBr USP (Cough Suppressant)

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose
- Cough

Alcohol 10%

Total 24 FL OZ (710 mL)

Two 12 FL OZ (355 mL) Bottles



# AUROQUIL COLD AND FLU NIGHTTIME RELIEF

acetaminophen, doxylamine succinate, and dextromethorphan hbr solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58602-137
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ANISE (UNII: 21C2F5E8RE)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL 1600 (UNII: 1212Z7S33A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

## Product Characteristics



<b>Color</b>	GREEN		<b>Score</b>	
<b>Shape</b>			<b>Size</b>	
<b>Flavor</b>	ANISE, MENTHOL		<b>Imprint Code</b>	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:58602-137-18	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/09/2015	10/01/2021
2	NDC:58602-137-20	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/09/2015	10/01/2021
3	NDC:58602-137-35	2 in 1 CARTON	02/09/2015	10/01/2021
3	NDC:58602-137-16	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL		part341	02/09/2015	10/01/2021

**Labeler** - Aurohealth LLC (078728447)

**Registrant** - Aurohealth LLC (078728447)

Establishment			
Name	Address	ID/FEI	Business Operations
Aurohealth LLC		078728447	MANUFACTURE(58602-137)